Two-loop additive mass renormalization with clover fermions and Symanzik improved gluons

A. Skouroupathis, M. Constantinou and H. Panagopoulos
Department of Physics, University of Cyprus, P.O. Box 20537, Nicosia CY-1678, Cyprus
email: php4as01@ucy.ac.cy, phpgmc1@ucy.ac.cy, haris@ucy.ac.cy
(February 2, 2008)

arXiv:0801.3146v1 [hep-lat] 21 Jan 2008

Abstract

We calculate the critical value of the hopping parameter, κ_c, in Lattice QCD, up to two loops in perturbation theory. We employ the Sheikholeslami-Wohlert (clover) improved action for fermions and the Symanzik improved gluon action with 4- and 6-link loops.

The quantity which we study is a typical case of a vacuum expectation value resulting in an additive renormalization; as such, it is characterized by a power (linear) divergence in the lattice spacing, and its calculation lies at the limits of applicability of perturbation theory.

Our results are polynomial in $c_{\rm SW}$ (clover parameter) and cover a wide range of values for the Symanzik coefficients c_i. The dependence on the number of colors N and the number of fermion flavors N_f is shown explicitly. In order to compare our results to non perturbative evaluations of κ_c coming from Monte Carlo simulations, we employ an improved perturbation theory method for improved actions.

Keywords: Lattice QCD, Lattice perturbation theory, Hopping parameter, Improved actions.

PACS numbers: 11.15.Ha, 12.38.Gc, 11.10.Gh, 12.38.Bx

I. INTRODUCTION

In the present work, we calculate the additive renormalization of the fermion mass in lattice QCD, using clover fermions and Symanzik improved gluons. The calculation is carried out up to two loops in perturbation theory and it is directly related to the determination of the critical value of the hopping parameter, κ_c.

The clover fermion action [1] (SW) succesfully reduces lattice discretization effects and approaches the continuum limit faster. This justifies the extensive usage of this action in Monte Carlo simulations in recent years. The coefficient $c_{\rm SW}$ appearing in this action is a free parameter for the current work and our results will be given as a polynomial in $c_{\rm SW}$.

Regarding gluon fields, we employ the Symanzik improved action [2], which also aims at minimizing finite lattice spacing effects. For the coefficients parameterizing the Symanzik action, we consider several choices of values which are frequently used in the literature.

The lattice discretization of fermions introduces some well known difficulties; demanding strict locality and absence of doublers leads to breaking of chiral symmetry. In order to recover this symmetry in the continuum limit one must set the renormalized fermion mass (m_R) equal to zero. To achieve this, the mass parameter $m_\circ$ appearing in the Lagrangian must approach a critical value m_c , which is nonzero due to additive renormalization.

The mass parameter $m_\circ$ is directly related to the hopping parameter κ used in simulations. Its critical value, κ_c, corresponds to chiral symmetry restoration:

$$\kappa_c = \frac{1}{2\, m_c\, a + 8\, r} \tag{1}$$

where a is the lattice spacing and r is the Wilson parameter. Using Eq.(1), the non-renormalized fermion mass is given by:

$$m_B \equiv m_\circ - m_c = \frac{1}{2\, a}\left(\frac{1}{\kappa} - \frac{1}{\kappa_c}\right) \tag{2}$$

Thus, in order to restore chiral symmetry one must consider the limit $m_\circ \to m_c$. This fact points to the necessity of an evaluation of m_c .

The perturbative value of m_c is also a necessary ingredient in higher-loop calculations of the multiplicative renormalization of operators (see, e.g., Ref. [3]). In mass independent schemes, such renormalizations are typically defined and calculated at zero renormalized mass, and this entails setting the value of the Lagrangian mass equal to m_c .

Previous studies of the hopping parameter and its critical value have appeared in the literature for Wilson fermions - Wilson gluons [4] and for clover fermions - Wilson gluons [5,6]. The procedure and notation in our work is the same as in the above references.

Our results for κ_c (and consequently for the critical fermion mass) depend on the number of colors (N) and on the number of fermion flavors (N_f). Besides that, there is an explicit dependence on the clover parameter $c_{\rm SW}$ which, as mentioned at the beginning, is kept as a free parameter. On the other hand, the dependence of the results on the choice of Symanzik coefficients cannot be given in closed form; instead, we present it in a list of Tables and Figures.

The rest of the paper is organized as follows: In Sec. II we formulate the problem, define the discretized actions, and describe our calculation of the necessary Feynman diagrams.

Sec. III is a presentation of our results. Finally, in Sec. IV we apply to our one- and two-loop results an improvement method, proposed by us [7–9]. This method resums a certain infinite class of subdiagrams, to all orders in perturbation theory, leading to an improved perturbative expansion. We end this section with a comparison of perturbative and non-perturbative results. Our findings are summarized in Sec. V.

II. FORMULATION OF THE PROBLEM

We begin with the Wilson formulation of the QCD action on the lattice, with N_f flavors of degenerate clover (SW) [1] fermions. In standard notation, it reads:

$$\begin{aligned} S_L = S_G &+ \sum_f \sum_x (4r + m_{\rm o}) \bar{\psi}_f(x) \psi_f(x) \\ &- \frac{1}{2} \sum_f \sum_{x,\,\mu} \Big[\bar{\psi}_f(x) \, (r - \gamma_\mu) \, U_{x,\,x+\mu} \psi_f(x+\mu) + \bar{\psi}_f(x+\mu) \, (r + \gamma_\mu) \, U_{x+\mu,\,x} \psi_f(x) \Big] \\ &+ \frac{i}{4} \, c_{\rm SW} \sum_f \sum_{x,\,\mu,\,\nu} \bar{\psi}_f(x) \sigma_{\mu\nu} \hat{F}_{\mu\nu}(x) \psi_f(x), \end{aligned} \tag{3}$$

$$\text{where :} \qquad \hat{F}_{\mu\nu} \equiv \frac{1}{8} \, (Q_{\mu\nu} - Q_{\nu\mu}) \tag{4}$$

$$\begin{aligned} \text{and :} \qquad Q_{\mu\nu} = &\, U_{x,\,x+\mu} U_{x+\mu,\,x+\mu+\nu} U_{x+\mu+\nu,\,x+\nu} U_{x+\nu,\,x} \\ &+ U_{x,\,x+\nu} U_{x+\nu,\,x+\nu-\mu} U_{x+\nu-\mu,\,x-\mu} U_{x-\mu,\,x} \\ &+ U_{x,\,x-\mu} U_{x-\mu,\,x-\mu-\nu} U_{x-\mu-\nu,\,x-\nu} U_{x-\nu,\,x} \\ &+ U_{x,\,x-\nu} U_{x-\nu,\,x-\nu+\mu} U_{x-\nu+\mu,\,x+\mu} U_{x+\mu,\,x} \end{aligned} \tag{5}$$

The clover coefficient $c_{\rm SW}$ is treated here as a free parameter. Particular choices of values for $c_{\rm SW}$ have been determined both perturbatively [1] and non-perturbatively [10], so as to minimize $\mathcal{O}(a)$ effects. The Wilson parameter r is set to $r = 1$ henceforth; f is a flavor index; $\sigma_{\mu\nu} = (i/2)[\gamma_\mu, \gamma_\nu]$. Powers of the lattice spacing a have been omitted and may be directly reinserted by dimensional counting.

Regarding gluons, we use the Symanzik improved gauge field action, involving Wilson loops with 4 and 6 links[1]:

$$\begin{aligned} S_G = \frac{2}{g^2} \Big[&c_0 \sum_{\rm plaquette} {\rm Re\,Tr} \, \{1 - U_{\rm plaquette}\} + c_1 \sum_{\rm rectangle} {\rm Re\,Tr} \, \{1 - U_{\rm rectangle}\} \\ &+ c_2 \sum_{\rm chair} {\rm Re\,Tr} \, \{1 - U_{\rm chair}\} + c_3 \sum_{\rm parallelogram} {\rm Re\,Tr} \, \{1 - U_{\rm parallelogram}\} \Big] \end{aligned} \tag{6}$$

(g is the bare coupling constant). The lowest order expansion of this action, leading to the gluon propagator, is

[1] 1×1 *plaquette*, 1×2 *rectangle*, 1×2 *chair* (bent rectangle), and $1 \times 1 \times 1$ *parallelogram* wrapped around an elementary 3-d cube.

$$S_{\rm G}^{(0)} = \frac{1}{2}\int_{-\pi/a}^{\pi/a}\frac{d^4k}{(2\pi)^4}\sum_{\mu\nu}A_\mu^a(k)\left[G_{\mu\nu}(k) - \frac{\xi}{\xi-1}\hat{k}_\mu\hat{k}_\nu\right]A_\nu^a(-k) \tag{7}$$

where ξ is the gauge fixing parameter (see Eq.(10)) and:

$$G_{\mu\nu}(k) = \hat{k}_\mu\hat{k}_\nu + \sum_\rho\left(\hat{k}_\rho^2\delta_{\mu\nu} - \hat{k}_\mu\hat{k}_\rho\delta_{\rho\nu}\right)d_{\mu\rho}, \qquad \hat{k}_\mu = \frac{2}{a}\sin\frac{ak_\mu}{2}\,, \quad \hat{k}^2 = \sum_\mu\hat{k}_\mu^2$$
$$d_{\mu\nu} = (1-\delta_{\mu\nu})\left[C_0 - C_1\,a^2\hat{k}^2 - C_2\,a^2(\hat{k}_\mu^2+\hat{k}_\nu^2)\right] \tag{8}$$

The coefficients C_i are related to c_i by

$$C_0 = c_0 + 8c_1 + 16c_2 + 8c_3\,, \;\; C_1 = c_2 + c_3\,, \;\; C_2 = c_1 - c_2 - c_3 \tag{9}$$

The Symanzik coefficients must satisfy: $c_0 + 8c_1 + 16c_2 + 8c_3 = 1$, in order to reach the correct classical continuum limit. Aside from this requirement, the values of c_i can be chosen arbitrarily; they are normally tuned in a way as to ensure $\mathcal{O}(a)$ improvement.

As always in perturbation theory, we must introduce an appropriate gauge-fixing term to the action; in terms of the gauge field $Q_\mu(x)$ [$U_{x,\,x+\mu} = \exp(i\,g\,Q_\mu(x))$], it reads:

$$S_{gf} = \frac{1}{1-\xi}\sum_{x,\mu,\nu}\mathrm{Tr}\,\{\Delta_\mu^-Q_\mu(x)\Delta_\nu^-Q_\nu(x)\}, \qquad \Delta_\mu^-Q_\nu(x) \equiv Q_\nu(x-\hat{\mu}) - Q_\nu(x). \tag{10}$$

Having to compute a gauge invariant quantity, we can, for convenience, choose to work either in the Feynman gauge ($\xi = 0$) or in the Landau gauge ($\xi = 1$). Covariant gauge fixing produces the following action for the ghost fields ω and $\overline{\omega}$

$$S_{gh} = 2\sum_x\sum_\mu\mathrm{Tr}\left\{(\Delta_\mu^+\omega(x))^\dagger\Big(\Delta_\mu^+\omega(x) + ig\,[Q_\mu(x),\omega(x)] + \frac{1}{2}ig\left[Q_\mu(x),\Delta_\mu^+\omega(x)\right]\right.$$
$$\left.-\frac{1}{12}g^2\left[Q_\mu(x),\left[Q_\mu(x),\Delta_\mu^+\omega(x)\right]\right] + \cdots\Big)\right\}, \qquad \Delta_\mu^+\omega(x) \equiv \omega(x+\hat{\mu}) - \omega(x). \tag{11}$$

Finally, the change of integration variables from links to vector fields yields a Jacobian that can be rewritten as the usual measure term S_m in the action:

$$S_m = \frac{1}{12}Ng^2\sum_x\sum_\mu\mathrm{Tr}\,\{Q_\mu(x)Q_\mu(x)\} + \cdots \tag{12}$$

In S_{gh} and S_m we have written out only terms relevant to our computation. The full action is: $S = S_L + S_{gf} + S_{gh} + S_m$.

The bare fermion mass m_B must be set to zero for chiral invariance in the classical continuum limit. Terms proportional to r in the action, as well as the clover terms, break chiral invariance. They vanish in the classical continuum limit; at the quantum level, they induce nonvanishing, flavor-independent fermion mass corrections. Numerical simulation algorithms usually employ the hopping parameter,

$$\kappa \equiv \frac{1}{2\,m_{\rm o}\,a + 8\,r} \tag{13}$$

as an adjustable input. Its critical value, at which chiral symmetry is restored, is thus $1/8r$ classically, but gets shifted by quantum effects.

The renormalized mass can be calculated in textbook fashion from the fermion self–energy. Denoting by $\Sigma^L(p, m_{\rm o}, g)$ the truncated, one particle irreducible fermion two-point function, we have for the fermion propagator:

$$S(p) = \Big[i \mathring{\not{p}} + m(p) - \Sigma^L(p, m_{\rm o}, g) \Big]^{-1} \tag{14}$$

$$\text{where :} \qquad \mathring{\not{p}} = \frac{1}{a} \sum_\mu \gamma_\mu \sin(a p^\mu), \quad m(p) = m_{\rm o} + \frac{2r}{a} \sum_\mu \sin^2(a p^\mu/2)$$

To restore the explicit breaking of chiral invariance, we require that the renormalized mass vanish:

$$S^{-1}(0)\Big|_{m_{\rm o} \to m_c} = 0 \qquad \Longrightarrow \qquad m_c = \Sigma^L(0, m_c\,, g) \tag{15}$$

The above is a recursive equation for m_c, which can be solved order by order in perturbation theory.

We denote by dm the additive mass renormalization of $m_\circ$: $m_B = m_\circ - dm$. In order to obtain a zero renormalized mass, we must require $m_B \to 0$, and thus $m_\circ \to dm$. Consequently,

$$m_c = dm = dm_{(1-\rm loop)} + dm_{(2-\rm loop)} \tag{16}$$

At tree level, $m_c = 0$.

Two diagrams contribute to $dm_{(1-\rm loop)}$, shown in Fig. 1. In these diagrams, the fermion mass must be set to its tree level value, $m_{\rm o} \to 0$.

Fig. 1. One-loop diagrams contributing to $dm_{(1-\rm loop)}$.
Wavy (solid) lines represent gluons (fermions).

The quantity $dm_{(2-\rm loop)}$ receives contributions from a total of 26 diagrams, shown in Fig. 2. Genuine two-loop diagrams must again be evaluated at $m_{\rm o} \to 0$; in addition, one must include to this order the one-loop diagram containing an $\mathcal{O}(g^2)$ mass counterterm (diagram 23).

Certain sets of diagrams, corresponding to one-loop renormalization of propagators, must be evaluated together in order to obtain an infrared convergent result: These are diagrams 7+8+9+10+11, 12+13, 14+15+16+17+18, 19+20, 21+22+23.



Fig. 2. Two-loop diagrams contributing to $dm_{(2-\mathrm{loop})}$. Wavy (solid, dotted) lines represent gluons (fermions, ghosts). Crosses denote vertices stemming from the measure part of the action; a solid circle is a fermion mass counterterm.

III. COMPUTATION AND RESULTS

Given that the dependence of m_c on the Symanzik coefficients c_i cannot be expressed in closed form, we chose certain sets of values for c_i , presented in Table I, which are in common use [11–16]: Plaquette, Symanzik (tree level improved), Tadpole Improved Lüscher-Weisz (TILW), Iwasaki and DBW2. Actually, since the gluon propagator contains only the combinations C_1 and C_2 (Eq.(9)), all results for m_c can be recast in terms of C_1, C_2 and one additional parameter, say, c_2; in this case the dependence on c_2 (at fixed C_1, C_2) is polynomial of second degree.

The contribution dm_l of the l^{th} one-loop diagram to dm, can be expressed as:

$$dm_l = \frac{(N^2-1)}{N}\, g^2 \cdot \sum_{i=0}^{2} c_{\rm SW}^i\, \varepsilon_l^{(i)} \tag{17}$$

where $\varepsilon_l^{(i)}$ are numerical one-loop integrals whose values depend on C_1, C_2. The dependence on $c_{\rm SW}$ is seen to be polynomial of degree 2 ($i = 0,\, 1,\, 2$).

The contribution to dm from two-loop diagrams that do not contain closed fermion loops, can be written in the form

$$dm_l = \frac{(N^2-1)}{N^2}\, g^4 \cdot \sum_{i,j,k} c_{\rm SW}^i\, N^j\, c_2^k\, e_l^{(i,j,k)} \tag{18}$$

where the index l runs over all contributing diagrams, $j = 0, 2$ and $k = 0,\, 1,\, 2$ (since up to two vertices from the gluon action may be present in a Feynman diagram). The dependence on $c_{\rm SW}$ is now polynomial of degree 4 ($i = 0, \cdots, 4$). The coefficients $e_l^{(i,j,k)}$ (as well as $\tilde{e}_l^{(i)}$ of Eq.(19) below) are two-loop numerical integrals; once again, they depend on C_1, C_2. Finally, the contribution to dm from two-loop diagrams containing a closed fermion loop, can be expressed as

$$dm_l = \frac{(N^2-1)}{N}\, N_f\, g^4 \cdot \sum_{i=0}^{4} c_{\rm SW}^i\, \tilde{e}_l^{(i)} \tag{19}$$

where the index l runs over diagrams 12-13, 19-20. Summing up the contributions of all diagrams, dm assumes the form

$$\begin{aligned} dm = \sum_l dm_l = \frac{(N^2-1)}{N}\, g^2 \cdot \sum_i c_{\rm SW}^i\, \varepsilon^{(i)} + \frac{(N^2-1)}{N^2}\, g^4 \cdot \sum_{i,j,k} c_{\rm SW}^i\, N^j\, c_2^k\, e^{(i,j,k)} \\ + \frac{(N^2-1)}{N}\, N_f\, g^4 \cdot \sum_i c_{\rm SW}^i\, \tilde{e}^{(i)} \end{aligned} \tag{20}$$

In the above, $\varepsilon^{(i)}$, $e^{(i,j,k)}$, $\tilde{e}^{(i)}$ are the sums over all contributing diagrams of the quantities: $\varepsilon_l^{(i)}$, $e_l^{(i,j,k)}$, $\tilde{e}_l^{(i)}$, respectively (cf. Eqs.(17,18,19)).

The coefficients $\varepsilon^{(i)}$ lead to the total contribution of one-loop diagrams. Their values are listed in Table II, for the ten sets of c_i values shown in Table I. Similarly, results for the coefficients $e^{(i,j,k)}$ and $\tilde{e}^{(i)}$ corresponding to the total contribution of two-loop diagrams, are presented in Tables III-VII.

In order to enable cross-checks and comparisons, numerical per-diagram values of the constants $\varepsilon_l^{(i)}$, $e_l^{(i,j,k)}$ and $\tilde{e}_l^{(i)}$ are presented in Tables VIII-XII, for the case of the Iwasaki action. For economy of space, several vanishing contributions to these constants have simply been omitted. A similar breakdown for other actions can be obtained from the authors upon request.

The total contribution of one-loop diagrams, for $N = 3$ can be written as a function of the clover parameter $c_{\rm SW}$. In the case of the Plaquette, Iwasaki, and DBW2 actions, we find, respectively:

$$dm_{(1-{\rm loop})}^{\rm Plaquette} = g^2 \Big(-0.434285489(1) + 0.1159547570(3)\, c_{\rm SW} + 0.0482553833(1)\, c_{\rm SW}^2 \Big) \tag{21}$$

$$dm_{(1-{\rm loop})}^{\rm Iwasaki} = g^2 \Big(-0.2201449497(1) + 0.0761203698(3)\, c_{\rm SW} + 0.0262264231(1)\, c_{\rm SW}^2 \Big) \tag{22}$$

$$dm_{(1-{\rm loop})}^{\rm DBW2} = g^2 \Big(-0.0972070995(5) + 0.0421775310(1)\, c_{\rm SW} + 0.01141359801(1)\, c_{\rm SW}^2 \Big) \tag{23}$$

A similar process can be followed for two-loop diagrams. In this case, we set $N = 3$, $c_2 = 0$ and we use three different values for the flavor number: $N_f = 0$, 2, 3. Thus, for the Plaquette, Iwasaki and DBW2 actions, the total contribution is, respectively:

$$N_f = 0: \quad dm^{\rm Plaquette}_{(2-{\rm loop})} = g^4 \Big(-0.1255626(2) + 0.0203001(2)\, c_{\rm SW} + 0.00108420(7)\, c^2_{\rm SW} - 0.00116538(2)\, c^3_{\rm SW} - 0.0000996725(1)\, c^4_{\rm SW} \Big) \tag{24}$$

$$N_f = 2: \quad dm^{\rm Plaquette}_{(2-{\rm loop})} = g^4 \Big(-0.1192361(2) + 0.0173870(2)\, c_{\rm SW} + 0.00836498(8)\, c^2_{\rm SW} - 0.00485727(3)\, c^3_{\rm SW} - 0.0011561947(4)\, c^4_{\rm SW} \Big) \tag{25}$$

$$N_f = 3: \quad dm^{\rm Plaquette}_{(2-{\rm loop})} = g^4 \Big(-0.1160729(2) + 0.0159305(2)\, c_{\rm SW} + 0.0120054(1)\, c^2_{\rm SW} - 0.00670321(3)\, c^3_{\rm SW} - 0.0016844558(6)\, c^4_{\rm SW} \Big) \tag{26}$$

$$N_f = 0: \quad dm^{\rm Iwasaki}_{(2-{\rm loop})} = g^4 \Big(-0.0099523(2) - 0.0024304(5)\, c_{\rm SW} - 0.00232855(4)\, c^2_{\rm SW} - 0.00032100(2)\, c^3_{\rm SW} - 0.0000419365(1)\, c^4_{\rm SW} \Big) \tag{27}$$

$$N_f = 2: \quad dm^{\rm Iwasaki}_{(2-{\rm loop})} = g^4 \Big(-0.0076299(2) - 0.0040731(5)\, c_{\rm SW} + 0.00102758(6)\, c^2_{\rm SW} - 0.00242924(3)\, c^3_{\rm SW} - 0.000457690(2)\, c^4_{\rm SW} \Big) \tag{28}$$

$$N_f = 3: \quad dm^{\rm Iwasaki}_{(2-{\rm loop})} = g^4 \Big(-0.0064687(2) - 0.0048944(5)\, c_{\rm SW} + 0.00270565(7)\, c^2_{\rm SW} - 0.00348335(3)\, c^3_{\rm SW} - 0.000665567(2)\, c^4_{\rm SW} \Big) \tag{29}$$

$$N_f = 0: \quad dm^{\rm DBW2}_{(2-{\rm loop})} = g^4 \Big(+0.005099(2) - 0.0053903(7)\, c_{\rm SW} - 0.0011157(1)\, c^2_{\rm SW} - 0.00004482(2)\, c^3_{\rm SW} - 0.0000111470(2)\, c^4_{\rm SW} \Big) \tag{30}$$

$$N_f = 2: \quad dm^{\rm DBW2}_{(2-{\rm loop})} = g^4 \Big(+0.005944(2) - 0.0061840(7)\, c_{\rm SW} + 0.0002046(2)\, c^2_{\rm SW} - 0.0010177(3)\, c^3_{\rm SW} - 0.000125065(3)\, c^4_{\rm SW} \Big) \tag{31}$$

$$N_f = 3: \quad dm^{\rm DBW2}_{(2-{\rm loop})} = g^4 \Big(+0.006366(2) - 0.0065809(7)\, c_{\rm SW} + 0.0008648(2)\, c^2_{\rm SW} - 0.0015042(4)\, c^3_{\rm SW} - 0.000182023(5)\, c^4_{\rm SW} \Big) \tag{32}$$

In Figs. 3, 4, and 5 we present the values of $dm_{(2-{\rm loop})}$ for $N_f = 0$, 2, 3, respectively; the results are shown for all choices of Symanzik actions which we have considered, as a function of $c_{\rm SW}$ ($N = 3$, $c_2 = 0$). In all cases, the dependence on $c_{\rm SW}$ is rather mild. One observes that $dm_{(2-{\rm loop})}$ is significantly smaller for all improved actions, as compared to the plaquette action; in particular, in the case of DBW2, $dm_{(2-{\rm loop})}$ is closest to zero and it vanishes exactly around $c_{\rm SW} = 1$.

Another feature of these results is that they change only slightly with N_f, especially in the range $c_{\rm SW} < 1.5$. This is due to the small contributions of diagrams with closed fermion loops (diagrams 12, 13, 19, 20). By the same token, in the case of nondegenerate flavors, $dm_{(2-{\rm loop})}$ is expected to depend only weakly on the mass of the virtual fermion.



Fig. 3. Total contribution of two-loop diagrams, for $N = 3$, $N_f = 0$ and $c_2 = 0$. Legends appear in the same top-to-bottom order as the corresponding lines.



Fig. 4. Total contribution of two-loop diagrams, for $N = 3$, $N_f = 2$ and $c_2 = 0$. Legends appear in the same top-to-bottom order as the corresponding lines.



Fig. 5. Total contribution of two-loop diagrams, for $N = 3$, $N_f = 3$ and $c_2 = 0$. Legends appear in the same top-to-bottom order as the corresponding lines.

IV. IMPROVED PERTURBATION THEORY

We now apply our method of improving perturbation theory [7–9], based on resummation of an infinite subset of tadpole diagrams, termed "cactus" diagrams. In Ref. [9] we show how this procedure can be applied to any action of the type we are considering here, and it provides a simple, gauge invariant way of dressing, to all orders, perturbative results at any given order (such as the one- and two-loop results of the present calculation). Some alternative ways of improving perturbation theory have been proposed in Refs. [17,18]. In a nutshell, our procedure involves replacing the original values of the Symanzik and clover coefficients by improved values, which are explicitly computed in [9]. Applying at first this method to one-loop diagrams, the improved ("dressed") value $dm^{\rm dr}$ of the critical mass ($N = 3$, $c_2 = 0$) can be written as:

$$dm^{\rm dr}_{(1-{\rm loop})} = \sum_{i=0}^{2} \varepsilon^{(i)}_{dr}\, c^i_{\rm SW} \tag{33}$$

In comparing with $\varepsilon^{(i)}$ of Eq. (20), the quantity $\varepsilon^{(i)}_{dr}$ is the result of one-loop Feynman diagrams with dressed values for the Symanzik parameters, and it has already been multiplied by $g^2\,(N^2-1)\,/N$. The dependence of $\varepsilon^{(i)}_{dr}$ on g is quite complicated now, and cannot be given in closed form; instead $\varepsilon^{(i)}_{dr}$ must be computed numerically for particular choices of g.

Listed in Table XIII are the results for $\varepsilon_{dr}^{(i)}$ along with the value of $\beta = 2N/g^2$ corresponding to each one of the 16 actions used in this calculation.

An attractive feature of this improvement procedure is that it can be applied also to higher loop perturbative results, with due care to avoid double counting of the cactus diagrams which were already included at one loop. Ideally, of course, one loop improvement should already be adequate enough, so as to obviate the need to consider higher loops; indeed, we find this to be the case and, consequently, we limit our discussion of two-loop improvement to only the plaquette action ($\beta = 5.29$, $N = 3$, $N_f = 2$), the Iwasaki action ($\beta = 1.95$, $N = 3$, $N_f = 2$) and the DBW2 action ($\beta = 0.87$ and $\beta = 1.04$, $N = 3$, $N_f = 2$). Using these values, the contribution to $dm^{\rm dr}_{(2-{\rm loop})}$ is a polynomial in $c_{\rm SW}$:

$$\begin{aligned} dm^{\rm dr}_{(2-{\rm loop}),\,{\rm plaquette}} = & -0.77398(8) + 0.16330(4)\, c_{\rm SW} + 0.06224534(1)\, c_{\rm SW}^2 \\ & -0.0044006(9)\, c_{\rm SW}^3 - 0.00073780(6)\, c_{\rm SW}^4 \end{aligned} \tag{34}$$

$$\begin{aligned} dm^{\rm dr}_{(2-{\rm loop}),\,{\rm Iwasaki}} = & -0.0813302(9) + 0.043030(3)\, c_{\rm SW} + 0.0308196(2)\, c_{\rm SW}^2 \\ & -0.00767090(8)\, c_{\rm SW}^3 - 0.001160923(1)\, c_{\rm SW}^4 \end{aligned} \tag{35}$$

$$\begin{aligned} dm^{\rm dr}_{(2-{\rm loop}),\,{\rm DBW2}(\beta=0.87)} = & -0.044906(1) + 0.029449(4)\, c_{\rm SW} + 0.0239522(2)\, c_{\rm SW}^2 \\ & -0.0082231(1)\, c_{\rm SW}^3 - 0.001218955(4)\, c_{\rm SW}^4 \end{aligned} \tag{36}$$

$$\begin{aligned} dm^{\rm dr}_{(2-{\rm loop}),\,{\rm DBW2}(\beta=1.04)} = & -0.031260(1) + 0.021793(2)\, c_{\rm SW} + 0.0188027(2)\, c_{\rm SW}^2 \\ & -0.00705284(9)\, c_{\rm SW}^3 - 0.001055657(1)\, c_{\rm SW}^4 \end{aligned} \tag{37}$$

The comparison between the total dressed contribution $dm^{\rm dr} = dm^{\rm dr}_{(1-{\rm loop})} + dm^{\rm dr}_{(2-{\rm loop})}$ and the unimproved contribution, dm, for the plaquette action is exhibited in Fig. 6, as a function of $c_{\rm SW}$. Similarly, $dm^{\rm dr}$ for the Iwasaki and the DBW2 actions is shown in Fig. 7 and Fig. 8, respectively.



Fig. 6. Improved and unimproved values of dm up to two loops, as a function of $c_{\rm SW}$, for the plaquette action ($\beta = 5.29$, $N = 3$, $N_f = 2$).



Fig. 7. Improved and unimproved values of dm up to two loops, as a function of $c_{\rm SW}$, for the Iwasaki action ($\beta = 1.95$, $N = 3$, $N_f = 2$).



Fig. 8. Improved and unimproved values of dm up to two loops, as a function of $c_{\rm SW}$, for the DBW2 action ($N = 3$, $N_f = 2$). We set $\beta = 0.87$ (solid lines) and $\beta = 1.04$ (dotted lines).

Finally, in Table XIV, we present a comparison of dressed and undressed results, for some commonly used values of β, N_f, $c_{\rm SW}$, and we also compare with available non perturbative estimates for κ_c [10,19–22]. We observe that improved perturbation theory, applied to one-loop results, already leads to a much better agreement with the non perturbative estimates.

V. DISCUSSION

To recapitulate, in this paper we have calculated the critical mass m_c, and the associated critical hopping parameter κ_c, up to two loops in perturbation theory, using the clover action for fermions and the Symanzik improved gluon action with 4- and 6-link loops. The perturbative value of m_c is a necessary ingredient in the higher-loop renormalization of operators, in mass independent schemes: Such renormalizations are typically defined and calculated at vanishing renormalized mass, which amounts to setting the Lagrangian mass equal to m_c.

In our calculations, we have chosen for the Symanzik coefficients c_i a wide range of values, which are most commonly used in numerical simulations. The dependence of our results on the number of colors N and the number of fermion flavors N_f is shown explicitly. The dependence on the clover parameter $c_{\rm SW}$ is in the form of a fourth degree polynomial whose coefficients we compute explicitly; it is expected, of course, that the most relevant values for $c_{\rm SW}$ are those optimized for $\mathcal{O}(a)$ improvement, either at tree level ($c_{\rm SW} = 1$), or at one loop [1], or non-perturbatively [10].

Since m_c is gauge invariant, we chose to calculate it in the Feynman gauge. The propagator appearing in Feynman diagrams is the inverse of a nondiagonal matrix; while this inverse

can be written down explicitly, it is more convenient, and more efficient in terms of CPU time, to perform the inversion numerically. Integrations over loop momenta were performed as momentum sums on lattices of finite size L, where typically $L \lesssim 40$; extrapolation to $L \to \infty$ introduces a systematic error, which we estimate quite accurately.

Our results for m_c are significantly closer to zero in the case of Symanzik improved actions, as compared to the plaquette action. In particular, the DBW2 action stands out among the rest, in that m_c vanishes exactly for a value of $c_{\rm SW}$ around 1. Thus, improved actions seem to bring us quite near the point of chiral symmetry restoration. The dependence of m_c on the number of flavors is seen to be very mild. This fact would also suggest that, in the case of nondegenerate flavors, m_c should depend only weakly on the mass of the virtual fermion.

Finally, we have made some comparisons among perturbative and non-perturbative results for κ_c. While these are expected to differ for a power divergent additive renormalization, such as the quantity under study, we nevertheless find a reasonable agreement. This agreement is further enhanced upon using an improved perturbative scheme, which entails resumming, to all orders in the coupling constant, a dominant subclass of tadpole diagrams. The method, originally proposed for the Plaquette action (see Ref. [7]), was extended in Ref. [9] to encompass all possible gluon actions made of closed Wilson loops, and can be applied at any given order in perturbation theory. As would be desirable, one-loop improvement is seen to be already adequate to give a reasonable agreement among perturbative and non-perturbative values. Indeed, our results for $\kappa^{\rm dr}_{1-{\rm loop}}$ are significally closer to the non-perturbative evaluations, as shown in Table XIV; in fact, the two-loop dressing procedure introduces no further improvement to the comparison.

Acknowledgments: This work is supported in part by the Research Promotion Foundation of Cyprus (Proposal Nr: ΕΝΤΑΞ/0504/11, ΕΝΙΣΧ/0505/45).

TABLES

TABLE I. Input parameters c_0, c_1, c_3 ($c_2 = 0$)

Action	c_0	c_1	c_3
Plaquette	1.0	0	0
Symanzik	1.6666667	-0.083333	0
TILW, $\beta c_0 = 8.60$	2.3168064	-0.151791	-0.0128098
TILW, $\beta c_0 = 8.45$	2.3460240	-0.154846	-0.0134070
TILW, $\beta c_0 = 8.30$	2.3869776	-0.159128	-0.0142442
TILW, $\beta c_0 = 8.20$	2.4127840	-0.161827	-0.0147710
TILW, $\beta c_0 = 8.10$	2.4465400	-0.165353	-0.0154645
TILW, $\beta c_0 = 8.00$	2.4891712	-0.169805	-0.0163414
Iwasaki	3.648	-0.331	0
DBW2	12.2688	-1.4086	0

TABLE II. Total contribution of one-loop diagrams

Action	$\varepsilon^{(0)}$	$\varepsilon^{(1)}$	$\varepsilon^{(2)}$
Plaquette	-0.1628570582(5)	0.0434830339(1)	0.01809576875(4)
Symanzik	-0.12805490528(8)	0.0378314931(2)	0.01476335801(5)
TILW (8.60)	-0.10821568768(4)	0.03408560232(6)	0.01265991972(4)
TILW (8.45)	-0.10749185625(3)	0.0339409375(1)	0.01258108895(1)
TILW (8.30)	-0.1064962872(3)	0.0337409869(2)	0.012472434543(4)
TILW (8.20)	-0.1058799831(2)	0.0336166372(1)	0.0124050416(1)
TILW (8.10)	-0.1050866191(1)	0.03345591621(5)	0.012318127134(5)
TILW (8.00)	-0.10410447893(3)	0.03325593631(8)	0.012210297749(7)
Iwasaki	-0.08255435613(4)	0.0285451387(1)	0.00983490867(5)
DBW2	-0.0364526623(2)	0.01581657412(5)	0.004280099253(2)

TABLE III. Total contribution of two-loop diagrams of order $\mathcal{O}(N^2,\, c_2^0)$

Action	$e^{(0,2,0)}$	$e^{(1,2,0)}$	$e^{(2,2,0)}$	$e^{(3,2,0)}$	$e^{(4,2,0)}$
Plaquette	-0.01753602(2)	0.00259963(2)	-0.000155894(8)	-0.000163242(2)	-0.00001721759(2)
Symanzik	-0.00810366(1)	0.00095046(2)	-0.000404510(9)	-0.000107348(2)	-0.00001275904(1)
TILW (8.60)	-0.00437013(7)	0.00019403(5)	-0.00045894(1)	-0.000078117(3)	-0.00001020820(1)
TILW (8.45)	-0.00425575(7)	0.00016978(6)	-0.00045962(1)	-0.000077102(3)	-0.00001011451(1)
TILW (8.30)	-0.00410086(7)	0.00013682(7)	-0.00046040(1)	-0.000075713(3)	-0.00000998564(1)
TILW (8.20)	-0.00400636(6)	0.00011666(8)	-0.00046080(1)	-0.000074857(3)	-0.00000990584(1)
TILW (8.10)	-0.00388630(6)	0.00009097(9)	-0.00046123(1)	-0.000073760(3)	-0.00000980314(1)
TILW (8.00)	-0.00374009(6)	0.00005958(9)	-0.000461601(9)	-0.000072410(3)	-0.00000967600(1)
Iwasaki	-0.00112957(2)	-0.00052964(6)	-0.000436966(5)	-0.000045009(3)	-0.00000682353(1)
DBW2	0.0008481(2)	-0.00085301(8)	-0.00018540(1)	-0.000006164(3)	-0.00000173502(3)

TABLE IV. Total contribution of two-loop diagrams of order $\mathcal{O}(N^0,\, c_2^0)$

Action	$e^{(0,0,0)}$	$e^{(1,0,0)}$	$e^{(2,0,0)}$	$e^{(3,0,0)}$	$e^{(4,0,0)}$
Plaquette	0.01656633(2)	-0.00055904(1)	0.002622771(7)	0.000158125(2)	0.00004282674(2)
Symanzik	0.00605656(1)	0.000935801(6)	0.002120980(9)	0.000104973(2)	0.00002971553(1)
TILW (8.60)	0.00202637(3)	0.00157890(3)	0.001790242(9)	0.000076167(2)	0.00002260669(1)
TILW (8.45)	0.00190729(3)	0.00159800(3)	0.001777415(9)	0.000075164(3)	0.00002235603(1)
TILW (8.30)	0.00174666(3)	0.00162375(2)	0.001759689(9)	0.000073791(3)	0.00002201243(1)
TILW (8.20)	0.00164901(3)	0.00163939(2)	0.001748661(9)	0.000072944(3)	0.00002180041(1)
TILW (8.10)	0.00152532(3)	0.00165917(2)	0.001734421(9)	0.000071859(3)	0.00002152826(1)
TILW (8.00)	0.00137535(4)	0.00168310(3)	0.00171671(1)	0.000070522(3)	0.00002119259(1)
Iwasaki	-0.00103022(1)	0.00203254(1)	0.001313076(3)	0.000043949(3)	0.00001423324(1)
DBW2	-0.0018961(2)	0.0016130(3)	0.000413397(9)	0.000005057(3)	0.00000307480(3)

TABLE V. Total contribution of two-loop diagrams containing closed fermion loops

Action	$\tilde{e}^{(0)}$	$\tilde{e}^{(1)}$	$\tilde{e}^{(2)}$	$\tilde{e}^{(3)}$	$\tilde{e}^{(4)}$
Plaquette	0.00118621(2)	-0.000546197(8)	0.001365146(9)	-0.000692228(3)	-0.00019809791(7)
Symanzik	0.00081496(1)	-0.000448276(6)	0.001041379(8)	-0.000574521(3)	-0.0001453370(2)
TILW (8.60)	0.00063643(1)	-0.000389464(5)	0.000857737(3)	-0.000500011(5)	-0.0001148491(1)
TILW (8.45)	0.00063033(1)	-0.000387269(5)	0.000851127(3)	-0.000497194(5)	-0.0001137544(1)
TILW (8.30)	0.00062198(1)	-0.000384243(5)	0.000842047(3)	-0.000493307(5)	-0.0001122515(1)
TILW (8.20)	0.00061684(1)	-0.000382366(5)	0.000836433(3)	-0.000490894(5)	-0.0001113227(1)
TILW (8.10)	0.00061025(1)	-0.000379946(5)	0.000829214(4)	-0.000487781(4)	-0.0001101288(1)
TILW (8.00)	0.00060214(1)	-0.000376945(5)	0.000820289(4)	-0.000483915(4)	-0.0001086536(1)
Iwasaki	0.00043546(1)	-0.00030800(1)	0.000629274(8)	-0.000395294(3)	-0.0000779538(3)
DBW2	0.00015833(3)	-0.00014883(4)	0.00024756(2)	-0.00018242(5)	-0.0000213595(6)

TABLE VI. Total contribution of two-loop diagrams containing the parameter c_2 (part 1)

Action	$e^{(0,0,1)}$	$e^{(1,0,1)}$	$e^{(2,0,1)}$	$e^{(0,2,1)}$	$e^{(1,2,1)}$
Plaquette	0.077167(3)	-0.019808(3)	-0.0085415(2)	-0.047102(4)	0.010439(3)
Symanzik	0.034929(2)	-0.010895(2)	-0.0041454(2)	-0.017940(2)	0.004491(2)
TILW (8.60)	0.020247(1)	-0.007117(2)	-0.0024559(1)	-0.008702(1)	0.002251(1)
TILW (8.45)	0.019816(1)	-0.006998(2)	-0.0024050(1)	-0.008448(1)	0.002185(1)
TILW (8.30)	0.019235(1)	-0.006835(2)	-0.0023362(1)	-0.0081078(6)	0.0020973(9)
TILW (8.20)	0.018881(1)	-0.006736(2)	-0.0022942(1)	-0.0079023(7)	0.002044(1)
TILW (8.10)	0.018433(1)	-0.006609(2)	-0.0022410(1)	-0.0076431(9)	0.0019761(8)
TILW (8.00)	0.017888(1)	-0.006454(2)	-0.0021762(1)	-0.0073300(6)	0.0018940(6)
Iwasaki	0.0087615(7)	-0.003656(1)	-0.00107856(8)	-0.0027484(4)	0.0006646(5)
DBW2	0.0007907(2)	-0.0004889(3)	-0.00008343(2)	0.0001308(2)	-0.0001587(3)

TABLE VII. Total contribution of two-loop diagrams containing the parameter c_2 (part 2)

Action	$e^{(2,2,1)}$	$e^{(3,2,1)}$	$e^{(0,2,2)}$	$e^{(1,2,2)}$	$e^{(2,2,2)}$
Plaquette	0.0039245(3)	-0.0000842143(1)	-0.09448252(9)	0.02755993(3)	0.010521016(1)
Symanzik	0.0014622(1)	-0.0000454986(1)	-0.03417549(2)	0.01248953(1)	0.0041047891(2)
TILW (8.60)	0.0006472(1)	-0.00002872341(6)	-0.017374635(6)	0.007205477(3)	0.0021218443(2)
TILW (8.45)	0.0006251(1)	-0.00002818123(6)	-0.016917713(6)	0.007049188(2)	0.0020666192(2)
TILW (8.30)	0.0005954(1)	-0.00002744385(5)	-0.016304614(5)	0.006838088(3)	0.0019924047(3)
TILW (8.20)	0.0005775(1)	-0.00002699223(5)	-0.015933835(5)	0.006709626(4)	0.0019474604(2)
TILW (8.10)	0.0005550(1)	-0.00002641646(5)	-0.015466270(5)	0.006546741(4)	0.0018907121(3)
TILW (8.00)	0.0005279(1)	-0.00002571231(5)	-0.014902324(4)	0.006348924(5)	0.0018221643(3)
Iwasaki	0.00015719(6)	-0.00001249281(2)	-0.00596123(2)	0.00295502(1)	0.0007286816(4)
DBW2	-0.00002436(1)	-0.00000050404(9)	-0.00028731(2)	0.00020317(4)	0.0000278810(8)

TABLE VIII. Contribution of one-loop diagrams, for the Iwasaki action

i	$\varepsilon_1^{(i)}$	$\varepsilon_2^{(i)}$
0	-0.05602636832(2)	-0.02652798781(3)
1	0	0.0285451387(1)
2	0	0.00983490867(5)

TABLE IX. Contribution of diagrams 3, 4, 6, for the Iwasaki action

i	j	k	$e_3^{(i,j,k)}$	$e_4^{(i,j,k)}$	$e_6^{(i,j,k)}$
0	0	0	-0.0003923686(9)	-0.000743134(3)	-0.0000714882(8)
0	2	0	0.0002615791(6)	0.000495422(2)	0.0000357441(4)
1	0	0	0	0.001900337(2)	0
1	2	0	0	0.0017774410(9)	0
2	0	0	0	-0.0010339720(2)	0
2	2	0	0	-0.001041123(1)	0.0002799238(4)

TABLE X. Contribution of diagrams 7-11, 14-18, 24, 26, for the Iwasaki action

i	j	k	$e_{7-11}^{(i,j,k)}$	$e_{14-18}^{(i,j,k)}$	$e_{24}^{(i,j,k)}$	$e_{26}^{(i,j,k)}$
0	0	0	0.00042802(1)	-0.000195263(2)	0	0
0	0	1	0.0057103(7)	0.0030512(2)	0	0
0	2	0	-0.00111995(2)	-0.00029748(1)	0	-0.000298742(2)
0	2	1	-0.0022472(3)	-0.0008718(2)	0	0.0003705893(7)
0	2	2	-0.00371263(2)	-0.00224859(1)	0	0
1	0	0	0	0.00064534(1)	0	0
1	0	1	0	-0.003656(1)	0	0
1	2	0	0	0.00011079(6)	-0.000144897(2)	0.000429899(1)
1	2	1	0	0.0006450(5)	0.000248682(4)	-0.00022905(1)
1	2	2	0	0.00295502(1)	0	0
2	0	0	0	-0.000000974(1)	0	0
2	0	1	0	-0.00107856(8)	0	0
2	2	0	0	0.000141960(3)	0.000042314(2)	0.0003303085(7)
2	2	1	0	0.00039546(6)	0.00002909398(7)	-0.000267364(2)
2	2	2	0	0.0007286816(4)	0	0
3	2	0	0	0	0	-0.000019835(1)
3	2	1	0	0	0	-0.00001249281(2)

TABLE XI. Contribution of diagrams 12, 13, 19, 20, for the Iwasaki action

i	$\tilde{e}_{12-13}^{(i)}$	$\tilde{e}_{19-20}^{(i)}$
0	0.000261920(6)	0.000173538(9)
1	-0.0000308339(1)	-0.00027717(1)
2	0.000370942(2)	0.000258332(8)
3	0	-0.000395294(3)
4	0	-0.0000779538(3)

TABLE XII. Contribution of diagrams 21-23, 25, 27, 28, for the Iwasaki action

i	j	k	$e_{21-23}^{(i,j,k)}$	$e_{25}^{(i,j,k)}$	$e_{27}^{(i,j,k)}$	$e_{28}^{(i,j,k)}$
0	0	0	0.000373419(3)	-0.000158621(4)	-0.000094848(3)	-0.0001759336(5)
0	2	0	-0.000373419(3)	0.000079311(2)	0	0.0000879668(3)
1	0	0	-0.000887295(1)	0.0001396819(4)	0.000045158(4)	0.0001893113(5)
1	2	0	0.000887295(1)	0.000085189(2)	0	-0.000120480(1)
2	0	0	0.000194437(1)	-0.0000319392(3)	0.000168506(2)	-0.0000509266(2)
2	2	0	-0.000194437(1)	-0.000005787(2)	0	0.0000098758(1)
3	0	0	0.000059183(3)	0	-0.000015234(1)	0
3	2	0	-0.000059183(3)	0.0000172022(5)	0	0.0000168072(6)
4	0	0	0.00000682353(1)	0	0.000007409712(6)	0
4	2	0	-0.00000682353(1)	0	0	0

TABLE XIII. Results for $dm^{\rm dr}_{(1-{\rm loop})}$ (Eq. (33)), with $N = 3$

Action	β	$\varepsilon^{(0)}_{dr}$	$\varepsilon^{(1)}_{dr}$	$\varepsilon^{(2)}_{dr}$
Plaquette	6.00	-0.579221119(2)	0.1159547570(3)	0.03618067788(9)
Symanzik	5.00	-0.4869797578(8)	0.1121369999(4)	0.03538605357(4)
Symanzik	5.07	-0.478756110(2)	0.11072412996(5)	0.03507238306(5)
Symanzik	6.00	-0.3915226522(2)	0.0947962001(5)	0.03124138429(9)
TILW (8.60)	3.7120	-0.5358770348(7)	0.1265917638(3)	0.03813963851(4)
TILW (8.45)	3.6018	-0.5497415338(3)	0.1291104644(3)	0.0386337113(1)
TILW (8.30)	3.4772	-0.5651407386(9)	0.1319263769(1)	0.0391695069(1)
TILW (8.20)	3.3985	-0.5756111531(9)	0.1337937558(7)	0.03951713046(7)
TILW (8.10)	3.3107	-0.5870122772(4)	0.1358437825(6)	0.0398899143(3)
TILW (8.00)	3.2139	-0.599415804(1)	0.138085996(2)	0.0402877133(4)
Iwasaki	1.95	-0.757856451(1)	0.1671007819(8)	0.044746728234(1)
Iwasaki	2.20	-0.6555102085(5)	0.1537748193(6)	0.04293183656(3)
Iwasaki	2.60	-0.541348980(1)	0.1359882440(3)	0.03967626495(6)
DBW2	0.6508	-0.7749943512(7)	0.1847244889(1)	0.04731717866(3)
DBW2	0.8700	-0.574781578(1)	0.1575688409(9)	0.04281261980(1)
DBW2	1.0400	-0.4822863343(9)	0.1412499230(5)	0.039186543574(5)

TABLE XIV. One- and two-loop results, and non-perturbative estimates for κ_c

Action	N_f	β	$c_{\rm SW}$	$\kappa_{1-{\rm loop}}$	$\kappa_{2-{\rm loop}}$	$\kappa^{\rm dr}_{1-{\rm loop}}$	$\kappa^{\rm dr}_{2-{\rm loop}}$	$\kappa_c^{\rm non-pert}$	[Ref.]
Plaquette	0	6.00	1.479	0.1301	0.1335	0.1362	0.1362	0.1392	[19]
Plaquette	0	6.00	1.769	0.1275	0.1306	0.1337	0.1332	0.1352	[10]
Plaquette	2	5.29	1.9192	0.1262	0.1307	0.1353	0.1341	0.1373	[20]
								0.1363	[21]
Iwasaki	2	1.95	1.53	0.1292	0.1368	0.1388	0.1379	0.1421	[22]
TILW (8.60)	0	3.7120	1.0	0.1339	0.1370	0.1378	0.1384		
TILW (8.00)	0	3.2139	1.0	0.1348	0.1387	0.1397	0.1406		
DBW2	2	0.87	0.0	0.1502	0.1384	0.1460	0.1479		
DBW2	2	0.87	1.0	0.1352	0.1372	0.1379	0.1379		
DBW2	2	1.04	0.0	0.1454	0.1375	0.1421	0.1434		
DBW2	2	1.04	1.0	0.1334	0.1348	0.1352	0.1352		

REFERENCES

[1] B. Sheikholeslami and R. Wohlert, Nucl. Phys. **B259**, 572 (1985).

[2] K. Symanzik, Nucl. Phys. **B226**, 187 (1983); K. Symanzik, Nucl. Phys. **B226**, 205 (1983).

[3] A. Skouroupathis and H. Panagopoulos, Phys. Rev. **D76**, 094514 (2007) [arXiv:0707.2906].

[4] E. Follana and H. Panagopoulos, Phys. Rev. **D63**, 017501 (2000) [hep-lat/0006001].

[5] H. Panagopoulos and Y. Proestos, Phys. Rev. **D65**, 014511 (2001) [hep-lat/0108021].

[6] S. Caracciolo, A. Pelissetto and A. Rago, Phys. Rev. **D64**, 094506 (2001) [hep-lat/0106013].

[7] H. Panagopoulos and E. Vicari, Phys. Rev. **D58**, 114501 (1998) [hep-lat/9806009].

[8] H. Panagopoulos and E. Vicari, Phys. Rev. **D59**, 057503 (1999) [hep-lat/9809007].

[9] M. Constantinou, H. Panagopoulos and A. Skouroupathis, Phys. Rev. **D74**, 074503 (2006) [hep-lat/0606001].

[10] M. Lüscher et al., Nucl. Phys. **B491**, 323 (1997) [hep-lat/9609035].

[11] M. Lüscher and P. Weisz, Phys. Lett. **B158**, 250 (1985).

[12] Y. Iwasaki, Univ. of Tsukuba Report UTHEP-118 (1983).

[13] K. Symanzik, Nucl. Phys. **B226**, 187 (1983).

[14] M. Lüscher and P. Weisz, Commun. Math. Phys. **97**, 59 (1985); Erratum-ibid. **98**, 433 (1985).

[15] M.G. Alford et al., Phys. Lett. **B361**, 87 (1995) [hep-lat/9507010].

[16] T. Takaishi, Phys. Rev. **D54**, 1050 (1996).

[17] G. Parisi, in: *Proceedings of the 20th International Conference on High Energy Physics*, Madison, 1980, ed. L. Durand and L. G. Pondrom (AIP, New York, 1981).

[18] G. P. Lepage and P. B. Mackenzie, Phys. Rev. **D48**, 2250 (1993) [hep-lat/9209022].

[19] UKQCD Collaboration (K.C. Bowler et al.), Phys. Rev. **D62**, 054506 (2000) [hep-lat/9910022].

[20] UKQCD Collaboration (C.R. Allton et al.), Phys. Rev. **D65**, 054502 (2002) [hep-lat/0107021].

[21] ALPHA Collaboration (M. Della Morte et al.), JHEP **0507**, 007 (2005) [hep-lat/0505026].

[22] CP-PACS Collaboration (A. Ali Khan et al.), Phys. Rev. **D65**, 054505 (2002); Erratum-ibid. **D67**, 059901 [hep-lat/0105015].